COVID-19 VACCINE PRIMARY SERIES AND BOOSTER ELIGIBILITY & TIMING

PFIZER-BIONTECH	MODERNA	JOHNSON & JOHNSON JANSSEN
Ages Authorized 5+ years old	Ages Authorized 18+ years old	Ages Authorized 18+ years old
Primary Series 2 doses, given 3 weeks (21 days) apart Moderately or severely immunocompromised people age 5 years and older should get an additional primary dose of Pfizer-BioNTech, given 28 days after 2nd shot.	Primary Series 2 doses, given 4 weeks (28 days) apart Moderately or severely immunocompromised people age 18 years and older should get an additional primary dose of Moderna, given 28 days after 2nd shot.	Primary Series 1 dose
Booster Dose (previously received Pfizer-BioNTech) Everyone ages 12 years and older, either Pfizer-BioNTech or Moderna (mRNA COVID-19 vaccines) at least 5 months after the last dose in their primary series. If receiving a Moderna booster, note that it is only authorized for age 18 years and older.	Booster Dose (previously received Moderna) Everyone ages 18 years and older, either Moderna or Pfizer-BioNTech (mRNA COVID-19 vaccines) at least 6 months after the last dose in their primary series.	Booster Dose (previously received J&J/Janssen) Everyone ages 18 years and older, either Pfizer-BioNTech or Moderna (mRNA COVID-19 vaccines) at least 2 months after the first dose of J&J/Janssen COVID-19 vaccine. Although not preferred by CDC, may get J&J/Janssen.
Up to Date on Vaccine Immediately after booster. If not eligible for booster yet, two weeks after completion of the primary series until eligible for booster.	Up to Date on Vaccine Immediately after booster. If not eligible for booster yet, two weeks after completion of the primary series until eligible for booster.	Up to Date on Vaccine Immediately after booster. If not eligible for booster yet, two weeks after completion of the primary series until eligible for booster.

In limited, exceptional situations where an individual received the first dose of an mRNA COVID-19 vaccine but is unable to complete the series with either the same or different mRNA COVID-19 vaccine (e.g., due to contraindication), a single dose of Janssen COVID-19 Vaccine may be considered at a minimum interval of 28 days from the mRNA COVID-19 vaccine dose if the person is ≥18 years of age. As part of a pre-vaccination discussion with a vaccination provider, all persons who elect to receive a Janssen COVID-19 Vaccine booster should be informed about the risk and symptoms of thrombosis with thrombocytopenia syndrome (TTS), and the need to seek immediate medical care should TTS symptoms develop. See the Contraindications and Precautions section for additional information on use of Janssen COVID-19 Vaccine and additional precautions in people with a contraindication to mRNA COVID-19 vaccines. People who receive Janssen COVID-19 Vaccine after a dose of an mRNA COVID-19 vaccine should be considered to have received a valid, single-dose Janssen vaccination - not a mixed primary vaccination series - and are considered fully vaccinated against COVID-19 ≥ 2 weeks after receipt of the single dose of the Janssen COVID-19 Vaccine.

Primary Series Source: <u>https://www.cdc.gov/coronavirus/2019-ncov/vaccines/different-vaccines.html</u>

Immunocompromised Source: https://www.cdc.gov/coronavirus/2019-ncov/vaccines/recommendations/immuno.html

Updated with ACIP 1/5/2022 meeting guidance, not yet published on CDC website.