

Clark County Youth Risk Behavior Survey (YRBS) 2025

2025 County-Wide High School Results

Introduction

The Youth Risk Behavior Survey (YRBS) is a survey that monitors health risk behaviors that contribute to the leading causes of death and disability among 9th-12th grade students. Topics include behaviors that contribute to unintentional injuries and violence, sexual behaviors, alcohol and drug use, tobacco use, dietary behaviors, physical activity, and the prevalence of obesity.

Methods

A census was conducted on youth risk behavior with 3,747 students across 14 high schools in Clark County. The overall response rate was 62.3%, which is considered a representative portion of the student population.

Participating Schools	
Catholic Central High School	School of Innovation
Global Impact STEM Academy	Shawnee High School
Greenon Senior High School	Southeastern High School
Kenton Ridge High School	Springfield High School
Nightingale Montessori High School	Springfield-Clark Career Technology Center
Northeastern High School	Springfield City Online
Northwestern Senior High School	Tecumseh High School

All incomplete or invalid responses were removed. Survey answers that were logically inconsistent with one another were also removed. Demographic variables including age, sex, race, and ethnicity were reported. The percentage (%) and 95% confidence interval were also reported for each question. Additionally, each response was analyzed by sex and race.

Data analysis was completed in collaboration with the Clark County Health Department by Alperen Korkmaz, PhD, Senior Statistical Consultant, Wright State University's Statistical Consulting Center under the Office of Research and Sponsored Programs (RSP).

Key Findings

Body Mass Index (BMI)

- 33.1% of students were overweight or obese.

Driving Habits

- 11.3% of students rode in a car driven by someone who had been drinking alcohol in the past 30 days.

Violence and Weapon Carrying

- 17.2% of students were involved in a physical fight during the past 12 months.
 - 65.3% of those students were male.
- 20.2% of students would be able to get a loaded gun without a parent or other adult's permission and be ready to fire within 10 minutes.
 - 63.4% of those students were male.

Bullying

- 21.5% of students were bullied on school property during the past 12 months.

- 11.4% of students have felt treated unfairly in school because of their race or ethnicity.

Depression, Suicide and Mental Health

- 12.7% of students have seriously considered attempting suicide in the past 12 months.
 - 67.0% of those students were female.
- 21.7% of students reported that their mental health was not good at least most of the time during the past 30 days.
- 30.5% of students felt sad or hopeless for two weeks in a row or more during the past 12 months.
- 32.6% of students rarely or never had a caring adult to go to about their feelings.

Tobacco

- 12.7% of students have tried smoking a cigarette.
- 22.9% of students have tried an electronic vapor product.
- 9.7% of students have smoked an electronic vapor product in the past 30 days.

Alcohol

- 33.4% of students have had at least one drink of alcohol 1 day or more in their life.
- 13.6% of students have had a drink of alcohol before the age of 13.

Marijuana

- 18.2% of students have used marijuana at least once in their life.
- 9.0% of students have used marijuana in the past 30 days.

Drugs

- 9.9% of students have taken non-prescribed pain medication or misused it at least once in their life.

Sexual Behavior

- 21.6% of students have had sexual intercourse.
- 11.5% of students used a condom when they last had sex.

Body Weight

- 35.3% of students self-describe as slightly or very overweight.

Physical Activity

- 15.9% of students have had a concussion at least once in the past 12 months from playing a sport or being physically active.

Other Topics

- 70.6% of students use social media several times a day or more often.
- 16.0% of students have at least sometimes gone hungry in the past 30 days because there was not enough food in their home.
- 26.0% of students have lived with a parent or guardian who was having a problem with alcohol or drug use.
- 11.8% of students rarely or never felt safe and secure in their neighborhood.
- 18.1% of students have gambled in the past 12 months.
 - 75.2% of those students were male.

Results

Population Demographics

Categories	Overall %	Female (49.9%)	Male (50.1%)
Age of Students			
12 or younger	0.2%	50.0%	50.0%
13 years	0.3%	45.5%	54.5%
14 years	19.1%	50.4%	49.6%
15 years	28.6%	50.6%	49.4%
16 years	25.3%	49.4%	50.6%
17 years	20.8%	50.1%	49.9%
18 or older	5.6%	46.6%	53.4%
Grade			
9th grade	29.3%	48.6%	51.4%
10th grade	27.8%	49.9%	50.1%
11th grade	25.3%	48.7%	51.3%
12th grade	17.4%	53.8%	46.2%
Ungraded or other	0.0%	0.0%	0.0%
Hispanic or Latino			
Yes	10.9%	50.0%	50.0%
No	89.1%	-	-
Race			
Multi-race	14.0%	49.8%	50.2%
American Indian or Alaska Native	1.4%	39.2%	60.8%
Asian	0.6%	50.0%	50.0%
Black or African American	11.3%	47.8%	52.2%
Native Hawaiian or Other Pacific Islander	0.3%	50.0%	50.0%
White	65.6%	50.2%	49.8%
Sex			
Female	49.9%	-	-
Male	50.1%	-	-
Sexual Orientation			
Heterosexual	75.9%	-	-
Gay or Lesbian	4.0%	-	-

Categories	Overall %	Female (49.9%)	Male (50.1%)
Bisexual	10.2%	-	-
Not Sure (Questioning)	2.9%	-	-
Other	2.9%	-	-
Gender Identity			
Transgender	2.9%	-	-
Not Transgender	92.4%	-	-
Unsure	2.0%	-	-

Body Mass Index (BMI)

Weight categories are defined by BMI percentiles. Please see Appendix I for definitions of weight categories by age group and gender.

Weight Categories	Overall %
Underweight	17.3%
Normal	49.7%
Overweight	18.0%
Obese	15.1%
Overweight & Obese	33.1%

Behavior Data

Questions	Overall % (95% CI)	Female (49.9%)	Male (50.1%)
Driving Habits			
Rarely or never wore a seat belt when riding in a car driven by someone else	6.3% (5.6% - 7.1%)	35.8%	64.2%
Rode in a car driven by someone who had been drinking alcohol, at least once in the past 30 days	11.3% (10.3% - 12.3%)	55.4%	44.6%
Texted or emailed while driving at least 1 day in the past 30 days	12.4% (11.4% - 13.5%)	46.5%	53.5%
Drove a car after they had been drinking alcohol at least 1 time in the past 30 days	2.0% (1.6% - 2.5%)	31.1%	68.9%
Violence and Weapon Carrying			
Carried a weapon (gun, knife, club) at least 1 day in the past 30 days on school property	1.9% (1.5% - 2.4%)	29.6%	70.4%
Ever seen someone get physically attacked, beaten, stabbed, or shot in their neighborhood	19.9% (18.7% - 21.2%)	48.6%	51.4%
Threatened or injured with a weapon on school property at least once in the past 12 months	4.7% (4.1% - 5.4%)	44.5%	55.5%
Carried a gun at least 1 day in the past 12 months	3.3% (2.7% - 3.9%)	20.7%	79.3%
Did not go to school because they felt unsafe at least 1 day in the past 30 days	8.7% (7.8% - 9.6%)	57.9%	42.1%
Involved in a physical fight at least once in the past 12 months	17.2% (16.0% - 18.5%)	34.7%	65.3%
Involved in a physical fight on school property at least once in the past 12 months	7.8% (6.9% - 8.7%)	34.8%	65.2%
Throughout life has at least sometimes been hit, beaten, kicked or physically hurt by an adult in the home	8.3% (7.4% - 9.2%)	62.7%	37.3%
Are able to get a loaded gun without a parent or adult's permission and be ready to fire within 10 minutes	20.2% (18.9% - 21.5%)	36.6%	63.4%
Sexual Violence			
Forced by anyone into doing sexual things at least once in the past 12 months	7.9% (7.0% - 8.8%)	76.6%	23.4%
Forced into having sexual intercourse	7.5% (6.7% - 8.4%)	73.7%	26.3%
Forced into doing sexual things by someone they were dating at least once in the past 12 months	6.1% (5.4% - 7.0%)	69.8%	30.2%
Physically hurt by someone they were dating at least once in the past 12 months	4.8% (4.2% - 5.6%)	52.6%	47.4%

Questions	Overall % (95% CI)	Female (49.9%)	Male (50.1%)
Bullying			
Bullied on school property in the past 12 months	21.5% (20.2% - 22.9%)	60.8%	39.2%
Electronically bullied in the past 12 months	14.3% (13.2% - 15.5%)	67.5%	32.5%
Have ever felt treated unfairly in school because of race or ethnicity	11.4% (10.5% - 12.5%)	57.2%	42.8%
Depression, Suicide and Mental Health			
Seriously considered attempting suicide in the past 12 months	12.7% (11.7% - 13.8%)	67.0%	33.0%
Made a plan to attempt suicide in the past 12 months	7.6% (6.8% - 8.5%)	70.3%	29.7%
Attempted suicide at least once in the past 12 months	4.6% (3.9% - 5.3%)	68.5%	31.5%
Had to be treated after a suicide attempt in the past 12 months	2.4% (2.0% - 3.0%)	55.2%	44.8%
Mental health was not good at least most of the time during the past 30 days	21.7% (20.4% - 23.0%)	69.4%	30.6%
Felt sad or hopeless for two weeks or more in a row in the past 12 months	30.5% (29.1% - 32.0%)	64.8%	35.2%
Rarely or never felt they had a caring adult to go to about their feelings	32.6% (31.1% - 34.1%)	50.6%	49.4%
Hurt themselves without wanting to kill themselves at least once in the past 12 months	15.0% (13.9% - 16.2%)	72.1%	27.9%
Tobacco and Electronic Vapor Products			
Tried smoking a cigarette	12.7% (11.6% - 13.8%)	48.2%	51.8%
Smoked a cigarette before age 13 years old	7.0% (6.2% - 7.9%)	49.0%	51.0%
Smoked cigarettes at least 1 day in the past 30 days	2.7% (2.2% - 3.3%)	39.6%	60.4%
Smoked cigarettes on at least 20 days in the past 30 days	0.7% (0.5% - 1.1%)	34.6%	65.4%
Smoked more than 10 cigarettes per day in the past 30 days	0.8% (0.6% - 1.2%)	34.5%	65.5%
Tried an electronic vapor product	22.9% (21.6% - 24.3%)	52.6%	47.4%

Questions	Overall % (95% CI)	Female (49.9%)	Male (50.1%)
Smoked an electronic vapor product at least once in the past 30 days	9.7% (8.8% - 10.7%)	52.3%	47.7%
Smoked an electronic vapor product 20 or more days in the past 30 days	3.7% (3.1% - 4.4%)	50.0%	50.0%
Got their electronic vapor products from a store (a vape shop, gas station, etc.) in the past 30 days	2.1% (1.7% - 2.6%)	43.4%	56.6%
Used electronic vapor products because they were feeling anxious, stressed, or depressed	5.1% (4.4% - 5.9%)	65.6%	34.4%
Used chewing tobacco, snuff, snus, etc. 1 day or more in the past 30 days	3.4% (2.9% - 4.1%)	30.2%	69.8%
Smoked cigars, cigarillos, or little cigars on 1 day or more in the past 30 days	3.2% (2.7% - 3.8%)	37.4%	62.6%
Alcohol			
Had at least one drink of alcohol 1 day or more in their life	33.4% (31.9% - 35.0%)	52.4%	47.6%
Had their first drink of alcohol before age 13 years old	13.6% (12.6% - 14.8%)	52.6%	47.4%
Had at least one drink of alcohol on at least 1 day in the past 30 days	10.6% (9.7% - 11.7%)	52.3%	47.7%
Usually got their alcohol from someone else in the past 30 days	5.7% (5.0% - 6.5%)	62.8%	37.2%
Had 5 drinks or more in a row in the past 30 days	4.5% (3.9% - 5.2%)	43.9%	56.1%
Had 4 drinks or more (female) or 5 drinks or more (male) on at least 1 day in the past 30 days	5.5% (4.9% - 6.3%)	46.5%	53.5%
Marijuana			
Had used marijuana at least once in their life	18.2% (17.0% - 19.5%)	48.9%	51.1%
Tried marijuana for the first time before age 13 years old	5.0% (4.3% - 5.7%)	48.6%	51.4%
Used marijuana at least once in the past 30 days	9.0% (8.1% - 10.0%)	49.2%	50.8%
Drugs			
Used a needle to inject an illegal drug at least once	1.6% (1.2% - 2.0%)	36.8%	63.2%
Have used any form of cocaine at least once in their life	1.8% (1.4% - 2.2%)	28.1%	71.9%

Questions	Overall % (95% CI)	Female (49.9%)	Male (50.1%)
Have sniffed glue, aerosols, paints to get high at least once in their life	4.1% (3.5% - 4.8%)	45.6%	54.4%
Have used heroin at least once in their life	1.4% (1.1% - 1.8%)	30.0%	70.0%
Have used methamphetamines at least once in their life	1.7% (1.3% - 2.2%)	32.3%	67.7%
Have used ecstasy at least once in their life	2.0% (1.6% - 2.5%)	32.9%	67.1%
Have taken non-prescribed pain medication or misused it at least once in their life	9.9% (9.0% - 10.9%)	59.1%	40.9%
Sexual Behavior			
Have had sexual intercourse	21.6% (20.3% - 22.9%)	49.2%	50.8%
First had sexual intercourse before age 13 years old	2.9% (2.4% - 3.5%)	43.4%	56.6%
Had sex with 4 or more people in their life	3.1% (2.6% - 3.7%)	41.8%	58.2%
Had sex with at least one person in the past 3 months	13.2% (12.1% - 14.3%)	48.6%	51.4%
Used alcohol or drugs the last time before sex	2.6% (2.1% - 3.1%)	46.2%	53.8%
Used a condom when they last had sex	11.5% (10.5% - 12.6%)	44.5%	55.5%
Did not use any method to prevent pregnancy when they last had sex	3.3% (2.8% - 3.9%)	45.8%	54.2%
Had sexual contact with both males and females	2.9% (2.4% - 3.5%)	72.9%	27.1%
Are non-heterosexual (LGBTQ+)	19.9% (18.7% - 21.2%)	73.9%	26.1%
Body Weight			
Self-describe as slightly or very overweight	35.3% (33.8% - 36.9%)	53.9%	46.1%
Not trying to do anything with their weight	17.1% (15.9% - 18.3%)	59.0%	41.0%
Dietary Behavior			
Did not eat fruit in the past 7 days	9.7% (8.8% - 10.7%)	42.7%	57.3%

Questions	Overall % (95% CI)	Female (49.9%)	Male (50.1%)
Did not eat green salad in the past 7 days	44.1% (42.5% - 45.7%)	44.2%	55.8%
Did not eat potatoes in the past 7 days	33.2% (31.7% - 34.8%)	51.9%	48.1%
Did not eat carrots in the past 7 days	52.6% (51.0% - 54.2%)	50.8%	49.2%
Did not eat other vegetables in the past 7 days	21.5% (20.2% - 22.8%)	46.1%	53.9%
Drank a soda or pop at least once per day in the past 7 days	21.5% (20.2% - 22.8%)	44.1%	55.9%
Did not eat breakfast on at least 1 day in the past 7 days	75.5% (74.1% - 76.8%)	53.3%	46.7%
Physical Activity			
Were active for 60 minutes per day less than 5 days in the past 7 days	49.6% (48.0% - 51.2%)	55.6%	44.4%
Did not go to Physical Education (PE) class in an average school week	75.8% (74.4% - 77.2%)	51.5%	48.5%
Did not play on any sports teams in the past 12 months	44.4% (42.8% - 46.0%)	49.8%	50.2%
Got a concussion from playing a sport or being active at least once in the past 12 months	15.9% (14.7% - 17.1%)	47.3%	52.7%
Other Health Topics			
Have ever been tested for HIV	4.5% (3.9% - 5.2%)	49.4%	50.6%
Have been tested for a STD other than HIV in the last 12 months	3.1% (2.6% - 3.7%)	48.6%	51.4%
Uses social media more than once a day	70.6% (69.1% - 72.1%)	52.8%	47.2%
Have not seen a dentist at least once in 12 months or more	24.4% (23.0% - 25.8%)	50.5%	49.5%
Have talked to a parent, guardian, or other trusted adult about STDs	36.6% (35.0% - 38.2%)	56.5%	43.5%
Get 4 hours or less of sleep on a school night	9.8% (8.9% - 10.8%)	50.4%	49.6%
Miscellaneous			
At least sometimes went hungry because there wasn't enough food at home in the last 30 days	16.0% (14.9% - 17.3%)	57.8%	42.2%

Questions	Overall % (95% CI)	Female (49.9%)	Male (50.1%)
Have gambled such as gambling on sports team, when playing cards or dice game, on the internet, lottery game or game of skill during the past 12 months	18.1% (16.9% - 19.4%)	24.8%	75.2%
Slept somewhere else besides their parent/guardians, family member or friends house in the past 30 days	3.5% (3.0% - 4.2%)	43.3%	56.7%
Did not have a usual place to sleep during the past 30 days	0.2% (0.1% - 0.5%)	12.5%	87.5%
Have lived with a parent or guardian who was having a problem with alcohol or drug use	26.0% (24.6% - 27.5%)	58.0%	42.0%
Rarely or never felt safe and secure in their neighborhood	11.8% (10.7% - 12.8%)	41.0%	59.0%

Behavior Data by Race

Questions	Overall %	Native Am. (1.4%)	Asian (0.6%)	Black/AA (11.3%)	Pacific Isl. (0.3%)	White (65.6%)	Multi-Race (14.0%)
Driving Habits							
Rarely or never wore a seat belt when riding in a car driven by someone else	6.3%	3.0%	0.4%	15.7%	0.9%	51.5%	23.0%
Rode in a car driven by someone who had been drinking alcohol, at least once in the past 30 days	11.3%	1.9%	0.5%	15.3%	0.7%	54.7%	17.5%
Texted or emailed while driving at least 1 day in the past 30 days	12.4%	1.1%	1.1%	9.1%	0.2%	66.6%	16.7%
Drove a car after they had been drinking alcohol at least 1 time in the past 30 days	2.0%	1.3%	2.7%	22.7%	1.3%	44.0%	20.0%
Violence and Weapon Carrying							
Carried a weapon (gun, knife, club) at least 1 day in the past 30 days on school property	1.9%	1.4%	2.8%	26.8%	2.8%	43.7%	15.5%
Ever seen someone get physically attacked, beaten, stabbed, or shot in their neighborhood	19.9%	1.8%	0.7%	12.2%	0.4%	58.5%	20.6%
Threatened or injured with a weapon on school property at least once in the past 12 months	4.7%	1.1%	1.1%	15.5%	1.1%	52.9%	21.3%
Carried a gun at least 1 day in the past 12 months	3.3%	3.3%	1.7%	19.0%	0.8%	51.2%	18.2%
Did not go to school because they felt unsafe at least 1 day in the past 30 days	8.7%	2.2%	1.6%	18.6%	0.6%	51.2%	16.8%
Involved in a physical fight at least once in the past 12 months	17.2%	3.0%	0.5%	14.1%	0.5%	55.8%	20.4%
Involved in a physical fight on school property at least once in the past 12 months	7.8%	3.1%	1.0%	15.3%	0.3%	53.0%	19.9%
Throughout life has at least sometimes been hit, beaten, kicked or physically hurt by an adult in the home	8.3%	2.0%	1.3%	11.4%	0.3%	57.5%	22.2%
Are able to get a loaded gun without a parent or adult's permission and be ready to fire within 10 minutes	20.2%	1.5%	0.4%	5.4%	0.1%	74.9%	13.6%

Questions	Overall %	Native Am. (1.4%)	Asian (0.6%)	Black/AA (11.3%)	Pacific Isl. (0.3%)	White (65.6%)	Multi-Race (14.0%)
Sexual Violence							
Forced by anyone into doing sexual things at least once in the past 12 months	7.9%	2.4%	0.7%	10.3%	1.0%	54.0%	23.7%
Forced into having sexual intercourse	7.5%	1.5%	0.4%	9.8%	1.1%	62.2%	20.4%
Forced into doing sexual things by someone they were dating at least once in the past 12 months	6.1%	3.1%	0.4%	11.0%	0.4%	58.6%	20.7%
Physically hurt by someone they were dating at least once in the past 12 months	4.8%	1.7%	1.1%	13.4%	1.1%	53.1%	22.3%
Bullying							
Bullied on school property in the past 12 months	21.5%	1.9%	0.4%	6.9%	0.5%	70.9%	15.8%
Electronically bullied in the past 12 months	14.3%	1.7%	0.4%	7.9%	0.6%	69.6%	15.7%
Have ever felt treated unfairly in school because of race or ethnicity	11.4%	1.2%	1.4%	26.9%	1.1%	27.6%	29.0%
Depression, Suicide and Mental Health							
Seriously considered attempting suicide in the past 12 months	12.7%	1.7%	0.9%	7.1%	0.4%	65.4%	19.2%
Made a plan to attempt suicide in the past 12 months	7.6%	1.8%	0.7%	7.3%	0.7%	68.1%	16.1%
Attempted suicide at least once in the past 12 months	4.6%	2.4%	0.6%	10.8%	0.6%	61.4%	17.5%
Had to be treated after a suicide attempt in the past 12 months	2.4%	2.2%	2.2%	13.5%	0.0%	48.3%	27.0%
Mental health was not good at least most of the time during the past 30 days	21.7%	1.6%	0.4%	8.7%	0.3%	67.7%	16.9%
Felt sad or hopeless for two weeks or more in a row in the past 12 months	30.5%	1.7%	0.4%	10.2%	0.4%	64.0%	17.4%
Rarely or never felt they had a caring adult to go to about their feelings	32.6%	1.3%	1.0%	15.4%	0.2%	56.7%	16.2%
Hurt themselves without wanting to kill themselves at least once in the past 12 months	15.0%	2.2%	1.3%	9.4%	0.2%	63.1%	18.0%

Questions	Overall %	Native Am. (1.4%)	Asian (0.6%)	Black/AA (11.3%)	Pacific Isl. (0.3%)	White (65.6%)	Multi-Race (14.0%)
Tobacco and Electronic Vapor Products							
Tried smoking a cigarette	12.7%	1.7%	0.4%	5.8%	0.2%	68.6%	17.5%
Smoked a cigarette before age 13 years old	7.0%	2.3%	0.8%	8.1%	0.4%	68.3%	15.4%
Smoked cigarettes at least 1 day in the past 30 days	2.7%	2.0%	2.0%	9.1%	0.0%	55.6%	23.2%
Smoked cigarettes on at least 20 days in the past 30 days	0.7%	3.7%	3.7%	11.1%	0.0%	37.0%	33.3%
Smoked more than 10 cigarettes per day in the past 30 days	0.8%	3.3%	3.3%	16.7%	0.0%	50.0%	23.3%
Tried an electronic vapor product	22.9%	1.8%	0.2%	8.9%	0.1%	63.6%	18.7%
Smoked an electronic vapor product at least once in the past 30 days	9.7%	1.4%	0.3%	8.5%	0.0%	63.7%	20.3%
Smoked an electronic vapor product 20 or more days in the past 30 days	3.7%	2.2%	0.0%	5.9%	0.0%	66.2%	20.6%
Got their electronic vapor products from a store (a vape shop, gas station, etc.) in the past 30 days	2.1%	1.3%	1.3%	11.5%	0.0%	60.3%	20.5%
Used electronic vapor products because they were feeling anxious, stressed, or depressed	5.1%	0.5%	0.5%	8.5%	0.0%	70.2%	16.0%
Used chewing tobacco, snuff, snus, etc. 1 day or more in the past 30 days	3.4%	0.8%	1.6%	11.8%	0.0%	62.2%	19.7%
Smoked cigars, cigarillos, or little cigars on 1 day or more in the past 30 days	3.2%	0.9%	0.9%	15.4%	0.0%	50.4%	24.8%
Alcohol							
Had at least one drink of alcohol 1 day or more in their life	33.4%	1.4%	0.5%	9.3%	0.3%	64.2%	16.7%
Had their first drink of alcohol before age 13 years old	13.6%	2.0%	0.8%	8.6%	0.2%	65.9%	15.5%
Had at least one drink of alcohol on at least 1 day in the past 30 days	10.6%	1.3%	0.3%	8.2%	0.0%	66.3%	20.1%
Usually got their alcohol from someone else in the past 30 days	5.7%	1.9%	0.5%	4.8%	0.0%	69.0%	20.0%
Had 5 drinks or more in a row in the past 30 days	4.5%	2.4%	0.6%	7.2%	0.0%	66.9%	18.7%

Questions	Overall %	Native Am. (1.4%)	Asian (0.6%)	Black/AA (11.3%)	Pacific Isl. (0.3%)	White (65.6%)	Multi-Race (14.0%)
Had 4 drinks or more (female) or 5 drinks or more (male) on at least 1 day in the past 30 days	5.5%	2.5%	0.5%	7.8%	0.0%	65.7%	18.1%
Marijuana							
Had used marijuana at least once in their life	18.2%	1.6%	0.4%	10.4%	0.1%	61.2%	21.4%
Tried marijuana for the first time before age 13 years old	5.0%	2.2%	1.1%	11.5%	0.5%	62.1%	18.1%
Used marijuana at least once in the past 30 days	9.0%	1.2%	0.6%	13.9%	0.0%	59.0%	21.7%
Drugs							
Used a needle to inject an illegal drug at least once	1.6%	3.4%	3.4%	24.1%	0.0%	34.5%	25.9%
Have used any form of cocaine at least once in their life	1.8%	3.1%	3.1%	24.6%	0.0%	41.5%	18.5%
Have sniffed glue, aerosols, paints to get high at least once in their life	4.1%	2.0%	2.0%	15.1%	0.0%	53.9%	21.7%
Have used heroin at least once in their life	1.4%	2.0%	2.0%	29.4%	0.0%	35.3%	21.6%
Have used methamphetamines at least once in their life	1.7%	3.2%	3.2%	23.8%	0.0%	39.7%	17.5%
Have used ecstasy at least once in their life	2.0%	1.4%	2.8%	20.8%	0.0%	48.6%	19.4%
Have taken non-prescribed pain medication or misused it at least once in their life	9.9%	1.4%	0.8%	15.6%	0.6%	56.2%	17.8%
Sexual Behavior							
Have had sexual intercourse	21.6%	1.5%	0.6%	11.8%	0.4%	62.7%	17.6%
First had sexual intercourse before age 13 years old	2.9%	2.8%	1.9%	14.8%	0.0%	59.3%	16.7%
Had sex with 4 or more people in their life	3.1%	1.8%	0.9%	16.7%	0.0%	50.9%	21.1%
Had sex with at least one person in the past 3 months	13.2%	0.8%	0.2%	10.4%	0.0%	66.0%	18.3%
Used alcohol or drugs the last time before sex	2.6%	2.1%	2.1%	13.8%	0.0%	56.4%	21.3%
Used a condom when they last had sex	11.5%	0.9%	0.0%	11.4%	0.0%	65.6%	14.9%
Did not use any method to prevent pregnancy when they last had sex	3.3%	1.7%	0.8%	20.7%	0.8%	50.4%	24.0%
Had sexual contact with both males and females	2.9%	2.8%	0.9%	5.6%	0.0%	62.0%	26.9%

Questions	Overall %	Native Am. (1.4%)	Asian (0.6%)	Black/AA (11.3%)	Pacific Isl. (0.3%)	White (65.6%)	Multi-Race (14.0%)
Are non-heterosexual (LGBTQ+)	19.9%	0.8%	0.8%	9.0%	0.4%	66.6%	17.9%
Body Weight							
Self-describe as slightly or very overweight	35.3%	1.1%	0.4%	7.5%	0.1%	65.9%	15.8%
Not trying to do anything with their weight	17.1%	0.8%	0.8%	11.3%	0.5%	69.6%	11.1%
Dietary Behavior							
Did not eat fruit in the past 7 days	9.7%	1.9%	0.8%	13.9%	0.6%	59.6%	18.7%
Did not eat green salad in the past 7 days	44.1%	1.4%	0.8%	11.6%	0.3%	65.1%	14.1%
Did not eat potatoes in the past 7 days	33.2%	1.6%	0.7%	11.8%	0.2%	65.5%	15.1%
Did not eat carrots in the past 7 days	52.6%	1.4%	0.7%	12.3%	0.2%	65.6%	14.0%
Did not eat other vegetables in the past 7 days	21.5%	1.5%	0.5%	12.5%	0.4%	64.1%	15.1%
Drank a soda or pop at least once per day in the past 7 days	21.5%	2.0%	0.6%	12.4%	0.2%	63.5%	15.1%
Did not eat breakfast on at least 1 day in the past 7 days	75.5%	1.4%	0.6%	10.4%	0.3%	66.0%	14.3%
Physical Activity							
Were active for 60 minutes per day less than 5 days in the past 7 days	49.6%	1.4%	0.7%	13.1%	0.3%	60.9%	15.5%
Did not go to Physical Education (PE) class in an average school week	75.8%	1.0%	0.6%	10.1%	0.2%	67.1%	14.2%
Did not play on any sports teams in the past 12 months	44.4%	1.7%	0.7%	12.9%	0.3%	60.3%	15.4%
Got a concussion from playing a sport or being active at least once in the past 12 months	15.9%	2.1%	0.9%	14.9%	0.4%	64.4%	13.5%
Other Health Topics							
Have ever been tested for HIV	4.5%	1.2%	0.6%	15.7%	1.2%	59.0%	17.5%
Have been tested for a STD other than HIV in the last 12 months	3.1%	1.8%	0.9%	14.2%	1.8%	51.3%	22.1%
Uses social media more than once a day	70.6%	1.2%	0.5%	11.1%	0.3%	65.1%	14.8%
Have not seen a dentist at least once in 12 months or more	24.4%	1.3%	0.8%	16.2%	0.3%	56.3%	16.1%

Questions	Overall %	Native Am. (1.4%)	Asian (0.6%)	Black/AA (11.3%)	Pacific Isl. (0.3%)	White (65.6%)	Multi-Race (14.0%)
Have talked to a parent, guardian, or other trusted adult about STDs	36.6%	0.8%	0.1%	9.1%	0.2%	68.9%	15.3%
Get 4 hours or less of sleep on a school night	9.8%	3.3%	1.1%	16.7%	0.8%	56.4%	16.1%
Miscellaneous							
At least sometimes went hungry because there wasn't enough food at home in the last 30 days	16.0%	2.4%	0.7%	15.3%	0.4%	56.1%	17.2%
Have gambled such as gambling on sports team, when playing cards or dice game, on the internet, lottery game or game of skill during the past 12 months	18.1%	2.4%	0.8%	11.6%	0.2%	62.5%	15.2%
Slept somewhere else besides their parent/guardians, family member or friends house in the past 30 days	3.5%	3.9%	0.8%	19.4%	0.8%	48.1%	19.4%
Did not have a usual place to sleep during the past 30 days	0.2%	0.0%	11.1%	22.2%	0.0%	11.1%	33.3%
Have lived with a parent or guardian who was having a problem with alcohol or drug use	26.0%	2.1%	0.7%	6.8%	0.2%	67.0%	18.2%
Rarely or never felt safe and secure in their neighborhood	11.8%	1.9%	0.9%	21.6%	0.4%	49.5%	15.6%

Appendix I: Weight Category Definitions

Weight categories are defined by BMI percentiles. To see the CDC weight category definitions for children and adolescents, please visit the CDC website at: https://www.cdc.gov/growthcharts/clinical_charts.htm